

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Problem Image Mailbox.**


# Flow Cytometric Analysis of HLA-A2+ Donor T Cells for HLA-A2 CMVpp65+ T Cells: Day 0 of Culture



Human PBMC were screened for HLA-A2 positivity. HLA-A2+ donors were screened with control (empty) HLA-A2 tetramers and CMVpp65 loaded tetramers. In the donor shown above, approximately 3% of the CD3+CD8+ express TCR specific for HLA-A2 CMVpp65.

FIG 1

## Flow Cytometric Analysis of CD25 Expression on HLA-A2 CMVpp65+ T Cells: Day 10 of Culture



PBMC were activated with CMV antigen (coated onto paramagnetic beads) and by day 10 of culture, many cells are shown to be CD25 (IL-2R) positive; and all of the HLA-A2 CMVpp65+ T cells are expressing high levels of CD25, indicating activation (right panel). Controls include the same donor cells treated with uncoated (antigen-negative) beads (middle panel), or an HLA-A2+ donor (donor 1) that did not show detectable tetramer+ cells at day 0 and was serologically negative for CMV (left panel). These data indicate that tetramer approaches can be effectively used to track antigen-specific T cells and their relative state of activation.

Title: COMPOSITIONS AND METHODS FOR ELIMINATING UNDESIRABLE SUBPOPULATIONS OF  
T CELLS IN PATIENTS WITH IMMUNOLOGICAL DEFECTS RELATED TO AUTOIMMUNITY  
AND ORGAN OR HEMATOPOIETIC STEM CELL TRANSPLANTATION

Inventors: Ronald Berenson et al.

Docket No.: 980034.422C1




FIG 3

Title: COMPOSITIONS AND METHODS FOR ELIMINATING UNDESIRABLE SUBPOPULATIONS OF T CELLS IN PATIENTS WITH IMMUNOLOGICAL DEFECTS RELATED TO AUTOIMMUNITY AND ORGAN OR HEMATOPOIETIC STEM CELL TRANSPLANTATION

Inventors: Ronald Berenson et al.

Docket No.: 980034.422C1

Mixing Xcellerated T Cells with Autologous B-CLL Leukemic Cells: Results in the Rapid Upregulation of Key Immunological Effector Molecules

Day 12 Xcellerated T Cells Co-cultured 24 hours with autologous leukemic B Cells




FIG 4

Title: COMPOSITIONS AND METHODS FOR ELIMINATING UNDESIRABLE SUBPOPULATIONS OF  
T CELLS IN PATIENTS WITH IMMUNOLOGICAL DEFECTS RELATED TO AUTOIMMUNITY  
AND ORGAN OR HEMATOPOIETIC STEM CELL TRANSPLANTATION

Inventors: Ronald Berenson et al.

Docket No.: 980034.422C1




FIG 5

# T Cells Grow and Tumor Cells Are Eliminated During the Xcellerate Process




FIG 6

Title: COMPOSITIONS AND METHODS FOR ELIMINATING UNDESIRE SUBPOPULATIONS OF  
T CELLS IN PATIENTS WITH IMMUNOLOGICAL DEFECTS RELATED TO AUTOIMMUNITY  
AND ORGAN OR HEMATOPOIETIC STEM CELL TRANSPLANTATION

Inventors: Ronald Berenson et al.

Docket No.: 980034.422C1




Fig. 7



Title: COMPOSITIONS AND METHODS FOR ELIMINATING UNDESIRE SUBPOPULATIONS OF  
T CELLS IN PATIENTS WITH IMMUNOLOGICAL DEFECTS RELATED TO AUTOIMMUNITY  
AND ORGAN OR HEMATOPOIETIC STEM CELL TRANSPLANTATION

Inventors: Ronald Berenson et al.

Docket No.: 980034.422C1

Xcellerate Process Restores Healthy T Cell Repertoire  
Rheumatoid Arthritis Patient




Fig. 8




Fig. 9




Fig. 10